



SOUTH ATLANTIC FISHERY MANAGEMENT COUNCIL

4055 Faber Place Drive, Suite 201, North Charleston SC 29405

Call: (843) 571-4366 | Toll-Free: (866) SAFMC-10 | Fax: (843) 769-4520 | Connect: www.safmc.net

Dr. Michelle Duval, Chair | Charlie Phillips, Vice Chair | Gregg T. Waugh, Executive Director

POLICY CONSIDERATIONS FOR SOUTH ATLANTIC FOOD WEBS AND CONNECTIVITY AND ESSENTIAL FISH HABITATS (December 2016)

Introduction

This document provides guidance from the South Atlantic Fishery Management Council (SAFMC) regarding South Atlantic Food Webs and Connectivity and the protection of Essential Fish Habitat (EFH) and Habitat Areas of Particular Concern (EFH-HAPCs) supporting the Council's move to Ecosystem Based Fishery Management. The guidance is consistent with the overall habitat protection policies of the SAFMC as formulated and adopted in the Habitat Plan (SAFMC 1998a), the Comprehensive EFH Amendment (SAFMC 1998b), the Fishery Ecosystem Plan of the South Atlantic Region (SAFMC 2009a), Comprehensive Ecosystem-Based Amendment 1 (SAFMC 2009b), Comprehensive Ecosystem-Based Amendment 2 (SAFMC 2011), Fishery Ecosystem Plan 2 (SAFMC, 2018), the Habitat Blueprint (SAFMC, 2023), and the various Fishery Management Plans (FMPs) of the Council.

For the purposes of policy, the findings assess potential threats and impacts to managed species EFH and EFH-HAPCs and the South Atlantic ecosystem associated with changes in food webs and connectivity and processes that could improve those resources or place them at risk. The policies and recommendations established in this document are designed to address such impacts in accordance with the habitat policies of the SAFMC as mandated by law. The SAMFC may revise this guidance in response to 1) changes in conditions in the South Atlantic region, 2) applicable laws and regulatory guidelines, 3) new knowledge about the impacts or 4) as deemed as appropriate by the Council.

For the sake of clarity, the following terms are to be defined as:

- Essential Fish Habitat (EFH): The aquatic habitat where fish spawn, breed, feed, and grow to maturity. This includes the physical, chemical, and biological properties of aquatic areas as well as their associated benthic habitats.
- Habitat Areas of Particular Concern (HAPC): subsets of EFH that are considered important to ecosystem function and are vulnerable to degradation or human impact e.g. spawning areas, nursery areas, pupping grounds, etc.

Policy Considerations

A key tenet of ecosystem-based fisheries management (EBFM) is the consideration of potential indirect effects of fisheries on food web linkages when developing harvest strategies and management plans. Examples of unintended consequences include the over exploitation of predators, an increase in abundance of their prey, and a decline of organisms two trophic levels below them, a phenomenon known as a trophic cascade (Carpenter et al. 1985). Alternatively, fishing on lower trophic level species, planktivorous "forage" fishes for example, may

ultimately lead to predator population declines due to food limitation (e.g. Okey et al. 2014; Walters and Martell 2004). Food web linkages connect different components of the larger ecosystem, such as pelagic forage fishes and their piscivorous predators or demersal carnivores. This connectivity between food webs over space, time, and depth creates multiple energy pathways that enhance ecosystem stability and resilience. Food web models are increasingly being utilized by fisheries managers as ecological prediction tools because they provide the capability to simulate the entire ecosystem from primary producers to top predators to fisheries. Food web models can serve to inform single species assessment and management and are capable of generating reference points (Walters et al. 2005) and ecosystem-level indicators (Coll et al. 2006; Fulton et al. 2005).




Figure 1-1. The marine food web of the South Atlantic, based on the latest iteration of the SA Ecopath with Ecosim model (SAFMC 2022). Nodes are colored based on type (see legend above). The pink nodes identify groups that include SAFMC managed species or are a managed species. The flow strength varies from Blue (weakest) to red (strongest). Each circle represents a group in the ecosystem and its size is logarithmically proportional to its biomass. Each line is a trophic interaction between two groups. The width of each line represents the flow proportion. The light grey horizontal lines indicate trophic levels. Diagram produced by Lauren Gentry, FWC-FWRI September 2024.

Threats to EFH and EFH-HAPCs from Changes in South Atlantic Food Web and Connectivity

The SAFMC finds that negative impacts to EFH and EFH-HAPCs can change South Atlantic food webs and connectivity for managed species. Table 1 following food webs and connectivity policy and research recommendations, presents a summary of South Atlantic fisheries and their designated EFH and EFH-HAPCs as presented in the SAFMC [EFH User Guide \(SAFMC 2024\)](#)

SAFMC Policies Addressing South Atlantic Food Webs and Connectivity

The SAFMC establishes the following policies to address South Atlantic food webs and connectivity, and to clarify and augment the general policies already adopted in the Habitat Plan and Comprehensive Habitat Amendment and Fishery Ecosystem Plan (SAFMC 1998a; SAFMC 1998b; SAFMC 2009a).

General Policies:

1. **Forage Fisheries**¹ - Managers should consider forage fish (Defined in footnote one as invertebrate and vertebrate species) stock abundances and dynamics, and their impacts on predator productivity, when setting catch limits to promote ecosystem sustainability. To do so, more science and monitoring information are needed to improve our understanding of the role of forage fish in the ecosystem. This information should be included in stock assessments, ecosystem models, and other fishery management tools and processes in order to support the development of sustainable harvest strategies that incorporate ecosystem considerations and trade-offs.
Note: Initial preliminary definition and potential list of forage species and forage fish species presented in Appendix A.
2. **Prey importance-** Diet data used to describe the importance of forage species in food web models may contain certain biases depending on the metric used to inform the relative contribution to the diet, and due to other challenges involved such as varying rates of digestion for different prey (Hyslop, 1980). Therefore, the evaluation of prey importance should consider a suite of appropriate indicators including mass, occurrence, and degree of overlap as a shared resource among multiple predators. Prey that are identified as important should be recognized as ecosystem component species, and considered within the appropriate fishery-management plan.
3. **Food Web Indicators** – Food web indicators have been employed to summarize the state of knowledge of an ecosystem or food web and could serve as ecological benchmarks to inform future actions.
4. **Food Web Connectivity** – Separate food webs exist in the South Atlantic, for example inshore-offshore, north-south, and benthic-pelagic, but they are connected by species that migrate between them such that loss of connectivity could have impacts on other components of the ecosystem that would otherwise appear unrelated

¹ NOAA defines Fishery as “Generally, a fishery is an activity leading to harvesting of fish. It may involve capture of wild fish or raising of fish through aquaculture. A unit determined by an authority or other entity that is engaged in raising or harvesting fish. Typically, the unit is defined in terms of some or all of the following: people involved, species or type of fish, area of water or seabed, method of fishing, class of boats, and purpose of the activities. The combination of fish and fishers in a region, the latter fishing for similar or the same species with similar or the same gear types”. and fish as a “Used as a collective term, includes mollusks, crustaceans and any aquatic animal which is harvested.”(United States, 2005)

and must be accounted for.

5. **Trophic Pathways** – Managers should aim to understand how fisheries production is driven either by bottom-up or top-down forcing and attempt to maintain diverse energy pathways to promote overall food web stability.
6. **Food Web Models** – Food web models can provide useful information to inform stock assessments, screen policy options for unintended consequences, examine ecological and economic trade-offs, and evaluate performance of management actions under alternative ecosystem states. A full Ecopath with Ecosim (EWE) model and a snapper grouper simplified model has been developed for the South Atlantic Region to be used as a tool for management decisions. (SAFMC, 2022)
7. **Ecosystem Component Species** – Ecosystem component species are species that do not require conservation and management under a federal fishery management plan, but are included in order to achieve ecosystem management objectives.
8. **Invasive Species** – Invasive species, most notably lionfish (*Pterois spp.*), are known to have negative effects on ecologically and economically important reef fish species through predation and competition and those effects should be accounted for in management actions.
9. **Contaminants** – Bioaccumulation of contaminants in food webs can have sub-lethal effects on marine fish, mammals, and birds and is also a concern for human seafood consumption.

Research and Information Needs Addressing South Atlantic Food Webs and Connectivity

1. Scientific research and collection of data to further understand the impacts of climate variability on the South Atlantic ecosystem and fish productivity must be prioritized. This includes research on species distribution, habitat, reproduction, recruitment, growth, survival, predator-prey interactions and vulnerability.
2. Characterization of offshore ocean habitats used by estuarine dependent species, which can be useful in developing ecosystem models.
3. Scientific research and monitoring to improve our understanding of the role of forage species in the ecosystem, in particular abundance dynamics and habitat use.
4. Basic data are the foundation of ecosystem-based fisheries management thus, fixing existing data gaps in the South Atlantic must be addressed first in order to build a successful framework for this approach in the South Atlantic.
5. NOAA in cooperation with regional partners should develop and evaluate an initial suite of products ([Climate Vulnerability Assessment](#); [Region Habitat Assessment](#)) at an ecosystem level to help prioritize the management and scientific needs in the South Atlantic region taking a systemic approach to identify overarching, common risks (i.e., climate change, rising sea level, changes in ocean current patterns and strength, etc.) across all habitat, taxa, ecosystem functions, fishery participants and dependent coastal communities.
6. NOAA in cooperation with regional partners should develop risk assessments to evaluate the vulnerability of South Atlantic species with respect to their exposure and

sensitivity to ecological and environmental factors affecting their populations.

7. NOAA should coordinate with ongoing regional modeling and management tool development efforts to ensure that ecosystem management strategy evaluations (MSEs) link to multispecies and single species MSEs, inclusive of economic, socio-cultural, and habitat conservation measures.
8. NOAA should develop ecosystem-level reference points (ELRPs) and thresholds as an important step to informing statutorily required reference points and identifying key dynamics, emergent ecosystem properties, or major ecosystem-wide issues that impact multiple species, stocks, and fisheries. Addressing basic data collection gaps is critical to successful development of ELRPs.
9. Continued support of South Atlantic efforts to refine EFH and HAPCs is essential to protect important ecological functions for multiple species and species groups in the face of climate change.

Habitats designated as EFH and EFH-HAPCs by the SAFMC (Table 1), if negatively impacted, can change South Atlantic food webs and connectivity for managed species.

Table 1. Habitats designated as Essential Fish Habitat (EFH), their associated managed fisheries/species, and EFH-HAPCs (Source: SAFMC EFH Users Guide 2024).

Essential Fish Habitat	Fisheries/Species	EFH- Habitat Areas of Particular Concern
Wetlands		
Estuarine and marine emergent wetlands	Shrimp, Snapper Grouper	Shrimp: State designated nursery habitats Mangrove wetlands
Tidal palustrine forested wetlands	Shrimp	
Submerged Aquatic Vegetation		
Estuarine and marine submerged aquatic vegetation	Shrimp, Snapper Grouper, Spiny lobster	Snapper Grouper, Shrimp
Shell bottom		
Oyster reefs and shell banks	Snapper Grouper	Snapper Grouper
Coral and Hardbottom		
Coral reefs, live/hardbottom, medium to high rock outcroppings from shore to at least 600 ft where the annual water temperature range is sufficient.	Snapper Grouper, Spiny lobster, Coral, Coral Reefs and Live Hard/bottom Habitat	The Point, Ten Fathom Ledge, Big Rock, MPAs; The <i>Phragmatopoma</i> (worm reefs) off central east coast of Florida and nearshore hardbottom; coral and hardbottom habitat from Jupiter through the Dry Tortugas, FL; Deepwater CHAPCs
rock overhangs, rock outcrops, manganese-phosphorite rock slab formations, and rocky reefs		Snapper-grouper [blueline tilefish]
Artificial reefs	Snapper Grouper	Special Management Zones
Soft bottom		
Subtidal, intertidal non-vegetated flats	Shrimp	
Offshore marine habitats used for spawning and growth to maturity	Shrimp	
Sandy shoals of capes and offshore bars	Coastal Migratory Pelagics	Sandy shoals; Capes Lookout, Fear, Hatteras, NC; Hurl Rocks, SC;
troughs and terraces intermingled with sand, mud, or shell hash at depths of 150 to 300 meters		Snapper-grouper [golden tilefish]
Water column		
Ocean-side waters, from the surf to the shelf break zone, including Sargassum	Coastal Migratory Pelagics	
All coastal inlets	Coastal Migratory Pelagics	Shrimp, Snapper-grouper
All state-designated nursery habitats of particular importance (e.g., PNA, SNA)	Coastal Migratory Pelagics	Shrimp, Snapper-grouper
High salinity bays, estuaries	Cobia in Coastal Migratory Pelagics	Spanish mackerel: Bogue Sound, New River, NC; Broad River, SC
Pelagic Sargassum	Dolphin	
Gulf Stream	Shrimp, Snapper-grouper, Coastal Migratory Pelagics, Spiny lobster, Dolphin-wahoo	
Spawning area in the water column above the adult habitat and the additional pelagic environment	Snapper-grouper	

Work Cited

- Carpenter, S.R., Kitchell, J F, Hodgson, J R. Bioscience 35.10 (1985): 634-639.
Cascading trophic interactions and lake productivity. Bioscience 35(10): 634- 639.
- Coll, M; Santojanni, A; Arneri, E; Palomera, I. 2006. An ecosystem model of the Northern and Central Adriatic Sea: analysis of ecosystem structure and fishing impacts. *Biologia marina mediterranea* 13.1: 467-471.
- Hyslop, E.J. 1980. Stomach content analysis-a review of methods and their application. *Journal of Fish Biology*. 17:411-429.
- Fulton, E., Fuller, M., Smith, A. and Punt, A. 2004. Ecological indicators of the ecosystem effects of fishing: final report. Australian Fisheries Management Authority Report R99/1546, pp. 116.
- Okey, T. A., A. M. Cisneros-Montemayor, R. Pugliese, and R. U. Sumaila. 2014. Exploring the trophodynamic signature of forage species in the U.S. South Atlantic Bight ecosystem. Fisheries Centre Working Paper 2014-14, University of British Columbia Fisheries Centre, Vancouver, Canada.
- SAFMC. 1998a. [Final Habitat Plan for the South Atlantic region: Essential Fish Habitat requirements for fishery management plans of the South Atlantic Fishery Management Council](#). South Atlantic Fishery Management Council, 1 Southpark Cir., Ste 306, Charleston, SC 29407-4699. 457 pp. plus appendices.
- SAFMC. 1998b. [Final Comprehensive Amendment Addressing Essential Fish Habitat in Fishery Management Plans of the South Atlantic Region. Including a Final Environmental Impact Statement /Supplemental Environmental Impact Statement, Initial Regulatory Flexibility Analysis, Regulatory Impact Review, and Social Impact Assessment/Fishery Impact Statement](#). South Atlantic Fishery Management Council, 1 Southpark Cir., Ste 306, Charleston, SC 29407-4699. 136pp.
- SAFMC (South Atlantic Fishery Management Council). 2009a. [Fishery Ecosystem Plan of the South Atlantic Region](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Ste 201, North Charleston, SC 29405.
- SAFMC (South Atlantic Fishery Management Council). 2009b. [Comprehensive Ecosystem-Based Amendment 1 for the South Atlantic Region](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Suite 201; North Charleston, SC 29405.
- SAFMC (South Atlantic Fishery Management Council). 2011. [Comprehensive Ecosystem-Based Amendment 2 for the South Atlantic Region](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Suite 201; North Charleston, SC 29405.

SAFMC (South Atlantic Fishery Management Council). 2018. [Fishery Ecosystem Plan II of the South Atlantic Region](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Ste 201, North Charleston, SC 29405.

SAFMC (South Atlantic Fishery Management Council). 2022. Ecosystem Model of Intermediate Complexity for Snapper Grouper Complex. South Atlantic Fishery Management Council, 4055 Faber Place Drive, Ste 201, North Charleston, SC 29405.

SAFMC (South Atlantic Fishery Management Council). 2023. [South Atlantic Fishery Management Council Habitat Program Evaluation and Blueprint](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Ste 201, North Charleston, SC 29405.

SAFMC (South Atlantic Fishery Management Council). 2024. [Users Guide to Essential Fish Habitat Designations by the South Atlantic Fishery Management Council](#). South Atlantic Fishery Management Council, 4055 Faber Place Drive, Ste 201, North Charleston, SC 29405.

Walters, Carl J, et. al. 2005, Possible ecosystem impacts of applying MSY policies from single-species assessment, ICES Journal of Marine Science, Volume 62, Issue 3, 2005, Pages 558–568, <https://doi.org/10.1016/j.icesjms.2004.12.005>

Walters and Mortell 2004. Fisheries Ecology And Management, Princeton University Press, Princeton, NJ 2004, ISBN 0-691-11545-1Paperback, 423 pp.

References

Andrews, S.N. et al. 2019. Consumption of Atlantic Salmon Smolt by Striped Bass: A Review of the Predator-Prey Encounter Literature and Implications for the Design of Effective Sampling Strategies. *Fishes* 2019, 4, 50; doi:10.3390/fishes4040050

Austin, Riley S. 2022. Age, Growth, Foraging, and Trophic Ecology of Bigeye (*Thunnus obesus*) and Yellowfin (*Thunnus albacares*) Tuna in Continental Shelf and Slope Regions of the Northeast U.S. MS Thesis, University of Maine. Electronic Theses and Dissertations. 3556.

Buchheister, A. et al. 2016. Spatial and temporal dynamics of Atlantic menhaden (*Brevoortia tyrannus*) recruitment in the Northwest Atlantic Ocean. *ICES Journal of Marine Science* (2016), 73(4), 1147– 1159. doi:10.1093/icesjms/fsv260

Grežlik, m.t. 2021. Evaluating the effects of Atlantic menhaden management and environmental change on the northwest Atlantic ocean ecosystem. Ms thesis, Humboldt state university. 91 pp.

Krumsick, K.J. 2020. Trophic and size spectra modeling reveal key species interactions and quantify community recovery dynamics within newfoundland and labrador fisheries ecosystems. PhD dissertation, Memorial University of Newfoundland. 250 pp.

Krumsick, K.J. and J.A.D. Fisher. 2022. Spatial variation in food web structure in a recovering marine ecosystem. *PLoS ONE* 17(5): e0268440. <https://doi.org/10.1371/journal.pone.0268440>

Luckhurst, B.E. 2018. A preliminary assessment of the ecological role and importance of

squid in the pelagic trophic web of the northwest atlantic ocean including the sargasso sea. Collect. Vol. Sci. Pap. Iccat, 74(7): 3679-3691.

- Murphy, KJ, GT Pecl, JD Everett, RF Heneghan, SA Richards, AJ Richardson, JM Semmens and JL Blanchard. 2023 Improving the biological realism of predator–prey size relationships in food web models alters ecosystem dynamics. Biol. Lett. 19: 20230142. <https://doi.org/10.1098/rsbl.2023.0142>
- Nadeau, S. 2021. Evaluating the Foraging Ecology and Energetics of Atlantic Bluefin Tuna (*Thunnus thynnus*) in the Gulf of Maine. MS Thesis, University of Maine. 155 pp. Electronic Theses and Dissertations. 3534. <https://digitalcommons.library.umaine.edu/etd/3534>
- Staudinger, M.D., et al. 2020. The role of sand lances (*Ammodytes* sp.) in the Northwest Atlantic Ecosystem: A synthesis of current knowledge with implications for conservation and management. Fish and Fisheries 21:522–556. DOI: 10.1111/faf.12445
- Strom, J.F., et al. 2019. Ocean predation and mortality of adult Atlantic salmon. Nature: Scientific Reports | (2019) 9:7890 | <https://doi.org/10.1038/s41598-019-44041-5>
- Turcotte, F. et al. 2023. Atlantic bluefin tuna diet variability in the southern Gulf of St. Lawrence, Canada. Marine Environmental Research 187 (2023) 105949.
- Varela, J.W., et al. 2020. Feeding ecology of Atlantic bluefin tuna (*Thunnus thynnus*) in the Gulf of Saint Lawrence, Canada. Marine Environmental Research Volume 161, October 2020, 105087. <https://doi.org/10.1016/j.marenvres.2020.105087>
- United States, National Oceanic and Atmospheric Administration. "NOAA fisheries glossary" , 2005
- Varela, J.W., et al. 2020. Feeding ecology of Atlantic bluefin tuna (*Thunnus thynnus*) in the Gulf of Saint Lawrence, Canada. Marine Environmental Research Volume 161, October 2020, 105087. <https://doi.org/10.1016/j.marenvres.2020.105087>
- Wuenschel, Mark J., et al. 2024. Variation in energy density of northwest Atlantic forage species: Ontogenetic, seasonal, annual, and spatial patterns. Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science. 2024;16:e10287. DOI: 10.1002/mcf2.10287

Appendix A. Below are the results of one method used to determine the top ten prey items of the predator species in each SAFMC Fishery Management Plan (FMP). The “Average %” value was calculated by the percent of each prey item in the FMP’s species’ diets averaged across all diets available for the species included in the FMP. For the Coastal Migratory Pelagic, Dolphin Wahoo, and Snapper grouper FMPs the top ten species for only fish prey was also calculated.

Top 10 Prey Items: Coastal Migratory Pelagic FMP			
If in diet, what's the average % that this prey makes up of the diet?			
All prey	Average %	Fish prey	Average %
Anchovies	15.811616	Anchovies	15.81162
Herrings	15.225112	Herrings	15.22511
Mega-invertebrate predators	11.594671	Sardines	10.4898
Sardines	10.489803	Shad	6.821819
Shad	6.8218191	Halfbeaks	6.484679
Halfbeaks	6.4846795	Scads	5.981667
Scads	5.9816669	Benthic coastal invertivores	3.334491
Benthic coastal invertivores	3.3344907	Demersal coastal invertivores	2.63137
Rock shrimps	2.851265	Pelagic coastal piscivores	2.400054
Demersal coastal invertivores	2.6313696	Pelagic planktivores	1.910574

Dolphin Wahoo FMP			
If in diet, what's the average % that this prey makes up of the diet?			
All prey	Average %	Fish prey	Average %
Sardines	15.05794	Sardines	15.05794
Pelagic oceanic piscivores	14.30661	Pelagic oceanic piscivores	14.30661
Auxis mackerels	9.303513	Auxis mackerels	9.303513
Squids	8.651381	Dolphinfish	6.876115
Dolphinfish	6.876115	Anchovies	6.251662
Anchovies	6.251662	Herrings	4.774037
Mega-invertebrate predators	5.403678	Demersal coastal omnivores	4.531909
Herrings	4.774037	Pelagic coastal piscivores	3.927871
Demersal coastal omnivores	4.531909	Blue runner	2.574502
Penaeid shrimps	4.329967	Pelagic planktivores	2.24857

Snapper Grouper FMP			
If in diet, what's the average % that this prey makes up of the diet?			
All prey	Average %	Fish prey	Average %
Mega-invertebrate predators	15.85355	Benthic coastal invertivores	6.770908
Benthic coastal invertivores	6.770908	Demersal coastal omnivores	5.177469
Echinoderms and gastropods	5.841086	Herrings	4.873793
Encrusting fauna	5.409635	Other grunts	4.765458
Squids	5.23724	Benthic oceanic invertivores	3.125774
Demersal coastal omnivores	5.177469	Demersal coastal invertivores	1.653085
Herrings	4.873793	Benthic coastal piscivores	1.607205
Other grunts	4.765458	Scads	1.399969
Bivalves/Oysters	3.315191	Demersal coastal piscivores	1.177001
Benthic oceanic invertivores	3.125774	Other sciaenids	1.123529

Golden Crab FMP			
If in diet, what's the average % that this prey makes up of the diet?			
All prey	Average %	Fish prey	Average %
Bivalves/Oysters		30.3	
Dead carcasses		19.6	
Offshore infaunal crustaceans		11.4	
Offshore benthic detritus		9.09	
Octopods		9.01	
Benthic oceanic invertivores		5	
Echinoderms and gastropods		4.81	
Mega-invertebrate predators		3.36	
Squids		2.44	
Penaeid shrimps		2.44	

Shrimp FMP			
If in diet, what's the average % that this prey makes up of the diet?			
All prey	Average %	Fish prey	Average %
Small mobile epifauna		21.95	
Mega-invertebrate predators		10	
Deep-burrowing infauna		9.793223	
Bivalves/Oysters		9.75	
Offshore polychaetes		7.69844	
Encrusting fauna		7.598771	
Echinoderms and gastropods		6.79991	
Estuarine benthic detritus		5.348395	
Offshore benthic detritus		5.348395	
Microphytobenthos		4.998501	

Spiny Lobster FMP

If in diet, what's the average % that this prey makes up of the diet?

All prey	Average %	Fish prey	Average %
Mega-invertebrate predators	20		
Offshore infaunal crustaceans	20		
Encrusting fauna	16		
Echinoderms and gastropods	10		
Bivalves/Oysters	10		
Offshore polychaetes	10		
Microbial heterotrophs	5		
Benthic macroalgae	5		
Carnivorous zooplankton	2		
Other zooplankton	2		

Acknowledgement: Many thanks to FWC-FWRI and Lauren Gentry for helping update the language, providing the graphs and conducting analysis to help inform the HEAP's Food web Policy.