



### South Atlantic Red Snapper 2020 Recreational Season Summary

Prepared for the South Atlantic Fishery Management Council

September 2020

# South Atlantic Red Snapper Monitoring in Florida

Florida Fish and Wildlife Conservation Commission
Fish and Wildlife Research Institute
Beverly.Sauls@MyFWC.com







#### Fisheries Dependent Monitoring in Florida

Mode	Special Surveys in-season	Year-round monitoring
Private boat	<ul><li>Effort monitoring at inlets</li><li>Dockside angler interviews</li><li>Dockside biological sampling</li></ul>	<ul> <li>MRIP</li> <li>State Reef Fish Survey – New!</li> </ul>
Charter	<ul><li>Mail/telephone survey of federally permitted vessels</li><li>Dockside biological sampling</li></ul>	MRIP For-Hire Survey
Headboat	No state sampling	<ul> <li>At-sea monitoring of discards</li> <li>NOAA Southeast Headboat Survey</li> </ul>
Commercial	<ul> <li>Enhanced TIP sampling during season</li> </ul>	

#### 2020 Recreational Season

- 8 inlets monitored each day
  - Good weather for offshore boating
  - Heavy boat traffic each day of the season
- 24 intercept survey assignments
  - 712 red snapper fishing parties interviewed
  - 2,684 red snapper anglers interviewed
  - 1,832 red snapper reported harvested
- 823 Red Snapper sampled for biological data
  - Average length 569 mm FL (22.4 inches)
  - Average weight 3.8 kg (8.37 lbs)
  - Otoliths collected for 724 fish







- New! Expanded to Atlantic coast July 2020
- Free endorsement required for private boat anglers to fish for:
  - Red Snapper
  - Vermilion, Mutton and Yellowtail snappers
  - Red, Black and Gag groupers
  - Gray Triggerfish, Hogfish, Amberjacks (4 Seriola sp.)
- Year-round, monthly estimates
  - Effort
  - Landings
  - Discards
- Special surveys will continue to be conducted to provide precise landings during mini-seasons

The FWC's

### State Reef Fish Survey

There are two components to the State Reef Fish Survey:



#### Together, these surveys tell us:

How many recreational reef fish trips are taken each month, and the total number of fish that are harvested and released.



As an angler or spear fisher, your responses on these surveys help improve scientific data and management of these important fisheries.



#### Estimates

- Special Red Snapper Surveys
  - Preliminary landings estimates for private boat and charter modes are in progress and will be available for September Council Meeting
  - Final report early 2021
- State Reef Fish Survey
  - Preliminary July estimate available late September



#### North Carolina

 121 carcasses sampled for otoliths from the North Carolina Carcass Collection Program:

-838mm FL max; 361mm FL min; 450mm FL avg

 Weather was fishable during both weekends; anecdotal reports from Marine Patrol indicate high activity



#### South Carolina


• In South Carolina, nine carcass collection freezer sites were available along the coast with a total of 92 fish racks contributed from recreational fishermen.

 Most samples were collected on the last day of the season. The total was slightly lower than 2019 (112 carcasses).






#### Red Snapper 2020 Mini-Season




#### Red Snapper Reported


- 129 Red Snapper were reported this year during the 4-day season.
- Longest red snapper reported: 40 inches
- Heaviest red snapper: 35.2 pounds


### Release Treatment by Depth





2019 2020






Release Treatment

• 14% to 15% of anglers used a descending device to release red snapper and other species during the 2020 red snapper mini-season.

## Catch Distribution by State



Length-Weight Comparison with SEDAR 41

