

# South Atlantic 2014 and 2015 Landings

Michael Larkin, Ph.D LAPP/DM Branch SERO

SAFMC SSC Meeting

October 20-22, 2015



#### **Notes on Commercial Landings Data**

- Landings were summarized in October of 2015.
  Preliminary 2015 landings are from January 1 to October 1, 2015.
- 2014 landings come from dealer trip tickets. 2015 landings come from in-season monitoring of dealer reported landings. In August 2014 dealer reporting switched from bimonthly to weekly.
- Landings are compiled and provided by the Southeast Fisheries Science Center.
- Landings are assigned by fisher reported catch area rather than dealer location



## 2014 Commercial Landings

| Species Complex                | Landings<br>(lbs) | 2014 ACL  | Units | Percent of ACL | Closure<br>Date |
|--------------------------------|-------------------|-----------|-------|----------------|-----------------|
| Atlantic spadefish             | 26,026            | 35,108    | ww    | 74%            |                 |
| Bar Jack                       | 4,759             | 5,265     | ww    | 90%            |                 |
| Black Grouper                  | 67,415            | 96,844    | ww    | 70%            |                 |
| Blueline tilefish              | 143,974           | 112,207   | ww    | 128%           | 6/23            |
| Cobia                          | 125,651           | 125,712   | ww    | 99%            | 12/11           |
| Deepwater complex <sup>a</sup> | 43,296            | 60,371    | ww    | 72%            | 7/10            |
| Dolphin                        | 1,128,241         | 1,157,001 | ww    | 98%            |                 |
| Gag                            | 324,447           | 326,722   | gw    | 99%            | 11/21           |
| Golden tilefish (H & L)        | 163,936           | 135,324   | gw    | 121%           | 8/29            |
| Golden tilefish (LongL)        | 531,769           | 405,971   | gw    | 131%           | 3/5             |
| Gray triggerfish               | 259,937           | 272,880   | ww    | 95%            | 5/12            |
| Grunts <sup>b</sup>            | 104,560           | 218,539   | ww    | 48%            |                 |
| Hogfish                        | 37,624            | 49,469    | ww    | 76%            |                 |

<sup>\*</sup> **Deepwater complex**<sup>a</sup> - Yellowedge grp, silk snapper, misty grp, queen snapper, sand tilefish, black snapper, blackfin snapper



<sup>\*</sup> **Grunts**<sup>b</sup> - White grunt, margate, sailor's choice, tomtate

<sup>\*</sup> **Dolphin**<sup>c</sup> – landings are from North, Mid, and South Atlantic regions



# 2014 Commercial Landings

| Species Complex                    | Landings<br>(lbs) | 2014 ACL | Units | Percent of ACL | Closure<br>Date |
|------------------------------------|-------------------|----------|-------|----------------|-----------------|
| Jacks <sup>c</sup>                 | 229,837           | 189,422  | ww    | 121%           | 7/15            |
| Mutton snapper                     | 91,173            | 157,743  | ww    | 58%            |                 |
| Porgies <sup>d</sup>               | 116,021           | 36,348   | ww    | 319%           | 12/9            |
| Red Grouper                        | 148,914           | 343,200  | ww    | 43%            |                 |
| Red Porgy                          | 150,634           | 154,500  | ww    | 97%            |                 |
| Red Snapper                        | 55,780            | 50,994   | gw    | 109%           | 9/9             |
| Scamp                              | 174,353           | 333,100  | ww    | 52%            |                 |
| Shallow water grouper <sup>e</sup> | 15,863            | 49,776   | ww    | 32%            |                 |
| Snappers <sup>f</sup>              | 145,118           | 215,662  | ww    | 67%            |                 |
| Snowy grouper                      | 91,413            | 82,900   | gw    | 110%           | 7/25            |

- \* **Jacks**<sup>c</sup> Almaco jack, banded rudderfish, lesser amberjack
- \* Porgies<sup>d</sup> Jolthead, knobbed, saucereye, whitebone, scup
- \* Shallow water grouper Red hind, rock hind, coney, graysby, yellowfin gpr, yellowmouth gpr
- \* **Snappers**<sup>f</sup> Gray, lane, cubera, dog, mahogany





# 2014 Commercial Landings

| Species Complex                      | Landings<br>(lbs) | 2014 ACL  | Units | Percent of ACL | Closure<br>Date |
|--------------------------------------|-------------------|-----------|-------|----------------|-----------------|
| Vermilion snapper<br>Jan 1 to Jun 30 | 461,601           | 446,080   | ww    | 103%           | 4/9             |
| Vermilion snapper<br>Jul to Dec 31   | 454,427           | 446,080   | ww    | 102%           | 9/12            |
| Wahoo <sup>c</sup>                   | 60,230            | 70,543    | ww    | 85%            |                 |
| Yellowtail snapper                   | 1,209,929         | 1,596,510 | ww    | 76%            |                 |

Wahoo<sup>c</sup> – landings are from North, Mid, and South Atlantic regions



# 2013/14 Commercial Landings

| Species<br>Complex   | Fishing Year | Landings<br>(lbs) | ACL     | Units | Percent of ACL | Closure Date |
|----------------------|--------------|-------------------|---------|-------|----------------|--------------|
| Greater<br>Amberjack | May 1-Apr 30 | 851,494           | 800,163 | ww    | 106%           |              |
| Black Sea Bass       | Jun 1-May 31 | 750,716           | 780,020 | ww    | 96%            |              |

## 2013/14 Preliminary Commercial Landings

| Species<br>Complex  | Fishing Year | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|---------------------|--------------|-------------------|-----------|-------|----------------|--------------|
| King Mackerel       | Mar 1-Feb 28 | 1,147,790         | 3,880,000 | ww    | 29%            |              |
| Spanish<br>Mackerel | Mar 1-Feb 28 | 3,152,991         | 3,130,000 | ww    | 101%           |              |

<sup>\*</sup> Landings from both S. Atlantic and Mid-Atlantic



2015 Preliminary Commercial Landings

| Species Complex                | Landings<br>(lbs) | 2015 ACL  | Units | Percent of ACL | Closure<br>Date |
|--------------------------------|-------------------|-----------|-------|----------------|-----------------|
| Atlantic spadefish             | 13,415            | 150,552   | ww    | 9%             |                 |
| Bar Jack                       | 2,571             | 13,228    | ww    | 19%            |                 |
| Black Grouper                  | 75,569            | 96,844    | ww    | 78%            |                 |
| Divoling tilefich              | 76 772            | 112,207   | ww    | 68%            |                 |
| Blueline tilefish              | 76,773            | 17,841    | ww    | 430%           | 4/7             |
| Cobia NY to GA                 | 24,921            | 60,000    | ww    | 42%            |                 |
| Cobia East Florida             | 50,203            | 70,000    | ww    | 72%            |                 |
| Deepwater complex <sup>a</sup> | 57,629            | 131,634   | ww    | 44%            |                 |
| Dolphin <sup>c</sup>           | 1,141,290         | 1,157,001 | ww    | 99%            | 6/30            |
| Gag                            | 222,306           | 295,459   | gw    | 75%            |                 |
| Golden tilefish (H & L)        | 102,737           | 135,324   | gw    | 76%            |                 |
| Golden tilefish (LongL)        | 385,734           | 405,971   | gw    | 95%            | 2/19            |
| G. Trigger Jan-Jun             | 251,312           | 272,880   | ww    | 92%            | 5/8             |
| G. Trigger Jul-Dec             | 101,391           | 63,918    | ww    | 159%           | 9/8             |
| Grunts <sup>b</sup>            | 75,321            | 217,903   | ww    | 35%            |                 |

<sup>\*</sup> Deepwater complexa - Yellowedge grp, silk snapper, misty grp, queen snapper, sand tilefish, black snapper, blackfin s



<sup>\*</sup> Gruntsb - White grunt, margate, sailor's choice, tomtate

<sup>\*</sup> **Dolphin**<sup>c</sup> – landings are from North, Mid, and South Atlantic regions



#### 2015 Preliminary Commercial Landings

| Species Complex                    | Landings<br>(lbs) | 2015 ACL | Units | Percent of ACL | Closure<br>Date |
|------------------------------------|-------------------|----------|-------|----------------|-----------------|
| Hogfish                            | 26,243            | 49,469   | Ww    | 53%            |                 |
| Jacks <sup>c</sup>                 | 238,127           | 189,422  | ww    | 126%           | 6/23            |
| Mutton snapper                     | 84,869            | 157,743  | ww    | 54%            |                 |
| Porgies <sup>d</sup>               | 28,527            | 36,348   | ww    | 78%            |                 |
| Red Grouper                        | 91,055            | 343,200  | ww    | 27%            |                 |
| Red Porgy                          | 112,343           | 164,000  | ww    | 69%            |                 |
| Red Snapper                        | 0                 | 0        | gw    | 0%             |                 |
| Scamp                              | 115,899           | 219,375  | ww    | 53%            |                 |
| Shallow water grouper <sup>e</sup> | 13,672            | 55,542   | ww    | 25%            |                 |
| Snappers <sup>f</sup>              | 152,237           | 344,884  | ww    | 44%            |                 |
| Snowy grouper                      | 133,031           | 115,451  | gw    | 115%           | 9/22            |

<sup>\*</sup> Jacks<sup>c</sup> - Almaco jack, banded rudderfish, lesser amberjack



<sup>\*</sup> Porgies<sup>d</sup> Jolthead, knobbed, saucereye, whitebone, scup

<sup>\*</sup> Shallow water grouper - Red hind, rock hind, coney, graysby, yellowfin gpr, yellowmouth gpr

<sup>\*</sup> **Snappers**<sup>f</sup> - Gray, lane, cubera, dog, mahogany

## 2015 Preliminary Commercial Landings

| Species Complex                      | Landings<br>(lbs) | 2015 ACL  | Units | Percent of ACL | Closure<br>Date |
|--------------------------------------|-------------------|-----------|-------|----------------|-----------------|
| Vermilion snapper<br>Jan 1 to Jun 30 | 446,195           | 438,260   | ww    | 102%           | 4/15            |
| Vermilion snapper<br>Jul to Dec 31   | 418,921           | 438,260   | ww    | 96%            | 9/22            |
| Wahoo <sup>c</sup>                   | 48,262            | 70,543    | ww    | 68%            |                 |
| Yellowtail snapper                   | 1,460,931         | 1,596,510 | ww    | 92%            |                 |

Wahoo<sup>c</sup> – landings are from North, Mid, and South Atlantic regions





#### 2014/15 Preliminary Commercial Landings

| Species<br>Complex   | Fishing Year   | Landings<br>(lbs) | ACL     | Units | Percent of ACL | Closure Date |
|----------------------|----------------|-------------------|---------|-------|----------------|--------------|
| Greater<br>Amberjack | May 1-Feb 28*  | 690,732           | 769,387 | gw    | 90%            |              |
| Black Sea Bass       | June 1-Dec 31* | 220,302           | 780,020 | ww    | 28%            |              |

\*2014/15 Fishing seasons were modified in Regulatory Amendment 14 for black sea bass and greater amberjack



## 2014/15 Preliminary Commercial Landings

| Species<br>Complex  | Fishing Year | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|---------------------|--------------|-------------------|-----------|-------|----------------|--------------|
| King Mackerel       | Mar 1-Feb 28 | 1,368,583         | 3,880,000 | ww    | 35%            |              |
| Spanish<br>Mackerel | Mar 1-Feb 28 | 2,815,218         | 3,330,000 | ww    | 85%            |              |

<sup>\*</sup> Landings from both S. Atlantic and Mid-Atlantic

## Notes on Recreational Landings Data

- Landings were summarized in September of 2015. Preliminary 2015 landings are from January to June (waves 1-3). Southeast Fisheries Science Center provided the landings.
- Landings were summarized using MRIP or MRFSS (calibrated from MRIP) depending on how the ACL is calculated.
- MRIP/MRFSS landings are available 45 days after the end of a two month wave.
- Landings include Headboat landings
- Landings estimates were updated by NMFS SERO to be consistent with ACL monitoring (e.g., post-stratification of estimates).



# 2014 Recreational Landings

| Species Complex                | Landings<br>(lbs) | 2014 ACL   | Units   | Percent of ACL | Closure<br>Date |
|--------------------------------|-------------------|------------|---------|----------------|-----------------|
| Atlantic spadefish             | 702,011           | 154,352    | ww      | 455%           |                 |
| Bar Jack                       | 1,979             | 19,515     | ww      | 10%            |                 |
| Black Grouper                  | 57,031            | 165,750    | ww      | 34%            |                 |
| Blueline tilefish              | 95,712            | 111,893    | ww      | 86%            |                 |
| Cobia                          | 1,179,031         | 1,445,687  | ww      | 82%            |                 |
| Deepwater complex <sup>a</sup> | 14,489            | 19,313     | ww      | 75%            |                 |
| Dolphin                        | 5,227,644         | 14,187,845 | ww      | 37%            |                 |
| Gag                            | 169,447           | 340,060    | gw      | 50%            |                 |
| Golden tilefish                | 1,357             | 3,019      | numbers | 45%            | 6/7             |
| Gray triggerfish               | 431,516           | 353,638    | ww      | 122%           | 11/26           |
| Grunts <sup>b</sup>            | 354,543           | 588,113    | ww      | 60%            |                 |
| Hogfish                        | 111,803           | 85,355     | ww      | 131%           |                 |
| Jacks <sup>c</sup>             | 226,004           | 267,799    | ww      | 84%            |                 |

<sup>\*</sup> Deepwater complexa - Yellowedge grp, silk snapper, misty grp, queen snapper, sand tilefish, black snapper, blackfin snapper



<sup>\*</sup> Grunts<sup>b</sup> - White grunt, margate, sailor's choice, tomtate

<sup>\*</sup> Jacks<sup>c</sup> - Almaco jack, banded rudderfish, lesser amberjack

<sup>\*</sup> MRFSS landings for gag and black grouper were post-stratified to include landings from the Florida Keys

<sup>\*</sup> Dolphin landings are from the North, Mid, and South Atlantic regions



### 2014 Recreational Landings

| Species Complex                    | Landings<br>(lbs) | 2014 ACL  | Units   | Percent of ACL | Closure<br>Date |
|------------------------------------|-------------------|-----------|---------|----------------|-----------------|
| Mutton snapper                     | 518,320           | 768,857   | ww      | 67%            |                 |
| Porgies <sup>d</sup>               | 128,231           | 106,914   | ww      | 120%           | 9/17            |
| Red Grouper                        | 38,756            | 436,800   | ww      | 9%             |                 |
| Red Porgy                          | 59,788            | 154,500   | ww      | 39%            |                 |
| Red Snapper <sup>e</sup>           | 31,683            | 22,576    | numbers | 140%           |                 |
| Scamp                              | 70,157            | 176,688   | ww      | 40%            |                 |
| Shallow water grouper <sup>f</sup> | 11,940            | 46,656    | ww      | 26%            |                 |
| Snappers <sup>g</sup>              | 830,700           | 728,577   | ww      | 114%           |                 |
| Snowy grouper                      | 1,214             | 523       | numbers | 232%           | 6/7             |
| Vermilion snapper                  | 406,070           | 419,840   | ww      | 97%            |                 |
| Wahoo                              | 656,871           | 1,724,418 | ww      | 38%            |                 |
| Wreckfish                          | 0                 | 11,750    | ww      | 0%             |                 |
| Yellowtail snapper                 | 933,279           | 1,440,990 | ww      | 65%            |                 |

<sup>\*</sup> Porgies<sup>d</sup> - Jolthead, knobbed, saucereye, whitebone, scup

<sup>\*</sup> Red Snapper<sup>e</sup> - Open during 8 days in July

<sup>\*</sup> Shallow water grouperf - Red hind, rock hind, coney, graysby, yellowfin gpr, yellowmouth gpr

<sup>\*</sup> **Snappers**<sup>g</sup> - Gray, lane, cubera, dog, mahogany

<sup>\*</sup> MRFSS landings for mutton snapper and yellowtail snapper were post-stratified to include landings from the Florida Keys

<sup>\*</sup> Wahoo landings are from the North, Mid, and South Atlantic regions

# 2013/14 Recreational Landings

| Species<br>Complex   | Fishing Year | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|----------------------|--------------|-------------------|-----------|-------|----------------|--------------|
| Greater<br>Amberjack | May 1-Apr 30 | 811,880           | 1,167,837 | ww    | 70%            |              |
| Black Sea Bass       | Jun 1-May 31 | 585,357           | 1,033,980 | ww    | 57%            |              |

- \* Greater amberjack landings include post-stratified MRFSS landings from Florida Keys.
- Black sea bass landings exclude post-stratified MRFSS landings from N. of Cape Hatteras

#### 2013/14 Recreational Landings

| Species<br>Complex   | Fishing Year | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|----------------------|--------------|-------------------|-----------|-------|----------------|--------------|
| King Mackerel*       | Mar 1-Feb 28 | 1,004,441         | 6,580,000 | ww    | 15%            |              |
| Spanish<br>Mackerel* | Mar 1-Feb 28 | 1,864,168         | 2,560,000 | ww    | 73%            |              |

<sup>\*</sup> Landings from both South Atlantic and Mid-Atlantic



2015 Preliminary Recreational Landings

| Species Complex                | Landings<br>(lbs) | 2015 ACL   | Units   | Percent of ACL | Closure<br>Date |
|--------------------------------|-------------------|------------|---------|----------------|-----------------|
| Atlantic spadefish             | 16,629            | 154,352    | ww      | 11%            |                 |
| Bar Jack                       | 3,573             | 19,515     | ww      | 18%            |                 |
| Black Grouper                  | 69,435            | 165,750    | ww      | 42%            |                 |
| Blueline tilefish              | 47,838            | 17,791     | ww      | 269%           | 6/10            |
| Cobia NY to GA                 | 974,081           | 630,000    | ww      | 155%           |                 |
| Cobia East Florida             | 333,028           | 830,000    | ww      | 40%            |                 |
| Deepwater complex <sup>a</sup> | 7,444             | 38,644     | ww      | 19%            |                 |
| Dolphin                        | 3,083,961         | 14,187,845 | ww      | 22%            |                 |
| Gag                            | 24,122            | 310,023    | gw      | 8%             |                 |
| Golden tilefish                | 3,843             | 3,019      | numbers | 127%           | 8/11            |
| Gray triggerfish               | 140,266           | 353,638    | ww      | 40%            |                 |
| Grunts <sup>b</sup>            | 115,759           | 588,113    | ww      | 20%            |                 |
| Hogfish                        | 243,948           | 85,355     | ww      | 286%           | 8/24            |
| Jacks <sup>c</sup>             | 75,652            | 267,799    | ww      | 28%            |                 |

<sup>\*</sup> Deepwater complexa - Yellowedge grp, silk snapper, misty grp, queen snapper, sand tilefish, black snapper, blackfin snapper



<sup>\*</sup> Grunts<sup>b</sup> - White grunt, margate, sailor's choice, tomtate

<sup>\*</sup> Jacks<sup>c</sup> - Almaco jack, banded rudderfish, lesser amberjack

<sup>\*</sup> MRFSS landings for gag and black grouper were post-stratified to include landings from the Florida Keys

<sup>\*</sup> Dolphin landings are for the North, Mid, and South Atlantic regions.



### 2015 Preliminary Recreational Landings

| Species Complex                    | Landings<br>(lbs) | 2015 ACL  | Units   | Percent of ACL | Closure<br>Date |
|------------------------------------|-------------------|-----------|---------|----------------|-----------------|
| Mutton snapper                     | 500,323           | 768,857   | ww      | 65%            |                 |
| Porgies <sup>d</sup>               | 66,403            | 106,914   | ww      | 62%            |                 |
| Red Grouper                        | 9,175             | 436,800   | ww      | 2%             |                 |
| Red Porgy                          | 58,569            | 164,000   | ww      | 36%            |                 |
| Red Snapper <sup>e</sup>           | 0                 | 22,576    | numbers | 0%             |                 |
| Scamp                              | 14,877            | 176,688   | ww      | 8%             |                 |
| Shallow water grouper <sup>f</sup> | 6,187             | 46,656    | ww      | 13%            |                 |
| Snappers <sup>g</sup>              | 236,630           | 728,577   | ww      | 32%            |                 |
| Snowy grouper                      | 618               | 4,152     | numbers | 15%            | 9/1             |
| Vermilion snapper                  | 154,415           | 412,480   | ww      | 37%            |                 |
| Wahoo                              | 265,640           | 1,724,418 | ww      | 15%            |                 |
| Wreckfish                          | 0                 | 21,650    | ww      | 0%             |                 |
| Yellowtail snapper                 | 461,710           | 1,440,990 | ww      | 32%            |                 |

<sup>\*</sup> Porgies<sup>d</sup> - Jolthead, knobbed, saucereye, whitebone, scup

<sup>\*</sup> Red Snapper<sup>e</sup> - Open during 8 days in July

<sup>\*</sup> Shallow water grouperf - Red hind, rock hind, coney, graysby, yellowfin gpr, yellowmouth gpr

<sup>\*</sup> **Snappers**<sup>9</sup> - Gray, lane, cubera, dog, mahogany

<sup>\*</sup> MRFSS landings for mutton snapper and yellowtail snapper were post-stratified to include landings from the Florida Keys

<sup>\*</sup> Wahoo landings are from the North, Mid, and South Atlantic regions

## 2014/15 Preliminary Recreational Landings

| Species<br>Complex   | Fishing Year | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|----------------------|--------------|-------------------|-----------|-------|----------------|--------------|
| King Mackerel*       | Mar 1-Feb 28 | 1,305,500         | 6,580,000 | ww    | 20%            |              |
| Spanish<br>Mackerel* | Mar 1-Feb 28 | 1,013,125         | 2,727,000 | ww    | 37%            |              |

<sup>\*</sup> Landings from both South Atlantic and Mid-Atlantic

### 2014/15 Preliminary Recreational Landings

| Species<br>Complex    | Fishing Year   | Landings<br>(lbs) | ACL       | Units | Percent of ACL | Closure Date |
|-----------------------|----------------|-------------------|-----------|-------|----------------|--------------|
| Greater<br>Amberjack* | May 1-Feb 28** | 859,457           | 1,167,837 | ww    | 74%            |              |
| Black Sea<br>Bass***  | Jun 1-Mar 31** | 357,589           | 1,033,980 | ww    | 36%            |              |

<sup>\*</sup> Greater amberjack landings include post-stratified MRFSS landings from Florida Keys.

<sup>\*\*</sup>Fishing seasons modified in Reg. Amendment 14 for black sea bass and greater amberjack

<sup>\*\*\*</sup>Black sea bass landings exclude post-stratified MRFSS landings from N. of Cape Hatteras

# QUESTIONS?