

## **Appendix I. Essential Fish Habitat and Move to Ecosystem Based Management**

### **South Atlantic Fishery Management Council Habitat Conservation, Ecosystem Coordination and Collaboration**

The Council, using the Essential Fish Habitat Plan as the cornerstone, adopted a strategy to facilitate the move to an ecosystem-based approach to fisheries management in the region. This approach required a greater understanding of the South Atlantic ecosystem and the complex relationships among humans, marine life, and the environment including essential fish habitat. To accomplish this, a process was undertaken to facilitate the evolution of the Habitat Plan into a Fishery Ecosystem Plan (FEP), thereby providing a more comprehensive understanding of the biological, social, and economic impacts of management necessary to initiate the transition from single species management to ecosystem-based management in the region.

#### **Moving to Ecosystem-Based Management**

The Council adopted broad goals for Ecosystem-Based Management to include maintaining or improving ecosystem structure and function; maintaining or improving economic, social, and cultural benefits from resources; and maintaining or improving biological, economic, and cultural diversity. Development of a regional FEP (SAFMC 2009a) provided an opportunity to expand the scope of the original Council Habitat Plan and compile and review available habitat, biological, social, and economic fishery and resource information for fisheries in the South Atlantic ecosystem. The South Atlantic Council views habitat conservation as the core of the move to EBM in the region. Therefore, development of the FEP was a natural next step in the evolution and expands and significantly updates the SAFMC Habitat Plan (SAFMC 1998a) incorporating comprehensive details of all managed species (SAFMC, South Atlantic States, ASMFC, and NOAA Fisheries Highly Migratory Species and Protected Species) including their biology, food web dynamics, and economic and social characteristics of the fisheries and habitats essential to their survival. The FEP therefore serves as a source document and presents more complete and detailed information describing the South Atlantic ecosystem and the impact of fisheries on the environment. This FEP updated information on designated Essential Fish Habitat (EFH) and EFH-Habitat Areas of Particular Concern; expanded descriptions of biology and status of managed species; presented information that will support ecosystem considerations for managed species; and described the social and economic characteristics of the fisheries in the region. In addition, it expanded the discussion and description of existing research programs and needs to identify biological, social, and economic research needed to fully address ecosystem-based management in the region. It is anticipated that the FEP will provide a greater degree of guidance by fishery, habitat, or major ecosystem consideration of bycatch reduction, prey-predator interactions, maintaining biodiversity, and spatial management needs. This FEP serves as a living source document of biological, economic, and social information for all Fishery Management Plans (FMP). Future Environmental Assessments and Environmental Impact Statements associated with subsequent amendments to Council FMPs will draw from or cite by reference the FEP.

The Fishery Ecosystem Plan for the South Atlantic Region encompasses the following volume structure:

- FEP Volume I - Introduction and Overview of FEP for the South Atlantic Region
- FEP Volume II - South Atlantic Habitats and Species
- FEP Volume III - South Atlantic Human and Institutional Environment
- FEP Volume IV - Threats to South Atlantic Ecosystem and Recommendations
- FEP Volume V - South Atlantic Research Programs and Data Needs
- FEP Volume VI - References and Appendices

Comprehensive Ecosystem-Based Amendment (CE-BA) 1 (SAFMC 2009b) is supported by this FEP and updated EFH and EFH-HAPC information and addressed the Final EFH Rule (e.g., GIS presented for all EFH and EFH-HAPCs). Management actions implemented in CE-BA 1 established deepwater Coral HAPCs to protect what is thought to be the largest continuous distribution (>23,000 square miles) of pristine, deepwater coral ecosystems in the world.

The Fishery Ecosystem Plan, slated to be revised every 5 years, will again be the vehicle to update and refine information supporting designation and future review of EFH and EFH-HAPCs for managed species. Planning for the update is being conducted in cooperation with the Habitat Advisory Panel during the fall and winter of 2013 with initiation during 2014.

### **Ecosystem Approach to Deepwater Ecosystem Management**

The South Atlantic Council manages coral, coral reefs and live/hard bottom habitat, including deepwater corals, through the Fishery Management Plan for Coral, Coral Reefs and Live/Hard Bottom Habitat of the South Atlantic Region (Coral FMP). Mechanisms exist in the FMP, as amended, to further protect deepwater coral and live/hard bottom habitats. The SAFMC's Habitat and Environmental Protection Advisory Panel and Coral Advisory Panel have supported proactive efforts to identify and protect deepwater coral ecosystems in the South Atlantic region. Management actions in Comprehensive Ecosystem-Based Amendment (CE-BA 1) (SAFMC 2009b) established deepwater coral HAPCs (C- HAPCs) to protect what is thought to be the largest continuous distribution (>23,000 square miles) of pristine deepwater coral ecosystems in the world. In addition, CE-BA 1 established areas within the CHAPC, which provide for traditional fishing in limited areas, which do not impact deepwater coral habitat. CE-BA 1, supported by the FEP, also addressed non-regulatory updates for existing EFH and EFH- HAPC information and addressed the spatial requirements of the Final EFH Rule (i.e., GIS presented for all EFH and EFH-HAPCs). Actions in this amendment included modifications in the management of the following: octocorals; special management zones (SMZs) off the coast of South Carolina; and sea turtle release gear requirements for snapper grouper fishermen. The amendment also designated essential fish habitat (EFH) and EFH-Habitat Areas of Particular Concern (EFH-HAPCs).

CE-BA 2 established annual catch limits (ACL) for octocorals in the South Atlantic as well as modifying the Fishery Management Unit (FMU) for octocorals to remove octocorals off the coast of Florida from the FMU (SAFMC 2011). The amendment also limited the possession of

managed species in the SMZs off South Carolina to the recreational bag limit for snapper grouper and coastal migratory pelagic species; modified sea turtle release gear requirements for the snapper grouper fishery based upon freeboard height of vessels; amends Council fishery management plans (FMPs) to designate or modify EFH and EFH-HAPCs, including the FMP for Pelagic Sargassum Habitat; amended the Coral FMP to designate EFH for deepwater Coral HAPCs designated under CE-BA 1; and amended the Snapper Grouper FMP to designate EFH-HAPCs for golden and blueline tilefish and the deepwater Marine Protected Areas. The final rule was published in the federal register on December 30, 2011, and regulations became effective on January 30, 2012.

### **Building from a Habitat to an Ecosystem Network to Support the Evolution**

Starting with our Habitat and Environmental Protection Advisory Panel, the Council expanded and fostered a comprehensive Habitat network in our region to develop the Habitat Plan of the South Atlantic Region completed in 1998 to support the EFH rule. Building on the core regional collaborations, the Council facilitated an expansion to a Habitat and Ecosystem network to support development of the FEP and CE-BA as well as coordinate with partners on other regional efforts.

#### *Integrated Ocean Observing System (IOOS) and Southeast Coastal and Ocean Observing Regional Association (SECOORA)*

The Integrated Ocean Observing System (IOOS®) is a partnership among federal, regional, academic, and private sector parties that works to provide new tools and forecasts to improve safety, enhance the economy, and protect our environment. IOOS supplies critical information about our Nation's oceans, coasts, and Great Lakes. Scientists working to understand climate change, governments adapting to changes in the Arctic, municipalities monitoring local water quality, and industries affected by coastal and marine spatial planning all have the same need: reliable, timely, and sustained access to data and information that inform decision making. Improving access to key marine data and information supports several purposes. IOOS data sustain national defense, marine commerce, and navigation safety. Scientists use these data to issue weather, climate, and marine forecasts. IOOS data are also used to make decisions for energy siting and production, economic development, and ecosystem-based resource management. Emergency managers and health officials need IOOS information to make decisions about public safety. Teachers and government officials rely on IOOS data for public outreach, training, and education.

SECOORA is one of 11 Regional Associations established nationwide through the US IOOS whose primary source of funding is through a 5-year cooperative agreement titled "Coordinated Monitoring, Prediction, and Assessment to Support Decision-Makers Needs for Coastal and Ocean Data and Tools". However, SECOORA was recently awarded funding via a NOAA Regional Ocean Partnership grant through the Governors' South Atlantic Alliance. SECOORA is the regional solution to integrating coastal and ocean observing data in the Southeast United States to inform decision makers and the general public. The SECOORA region encompasses 4 states, over 42 million people, and spans the

coastal ocean from North Carolina to the west Coast of Florida and is creating customized products to address these thematic areas: Marine Operations; Coastal Hazards; Ecosystems, Water Quality, Living Marine Resources; and Climate Change. The Council is a voting member and Council staff was recently re-elected to serve on the Board of Directors for the Southeast Coastal Regional Ocean Observing Association (SECOORA) to guide and direct priority needs for observation and modeling to support fisheries oceanography and integration into stock assessments through SEDAR. Cooperation through SECOORA is envisioned to facilitate the following:

- Refining current or water column designations of EFH and EFH-HAPCs (e.g., Gulf Stream and Florida Current).
- Providing oceanographic models linking benthic, pelagic habitats, and food webs.
- Providing oceanographic input parameters for ecosystem models.
- Integration of OOS information into Fish Stock Assessment process in the SA region.
- Facilitating OOS system collection of fish and fishery data and other research necessary to support the Council's use of area-based management tools in the SA Region including but not limited to EFH, EFH-HAPCs, Marine Protected Areas, Deepwater Coral Habitat Areas of Particular Concern, Special Management Zones, and Allowable Gear Areas.
- Integration of OOS program capabilities and research Needs into the South Atlantic Fishery Ecosystem Plan.
- Collaboration with SECOORA to integrate OOS products with information included in the Council's Habitat and Ecosystem Web Services and Atlas to facilitate model and tool development.
- Expanding Map Services and the Regional Habitat and Ecosystem Atlas in cooperation with SECOORAs Web Services that will provide researchers access to data or products including those collected/developed by SA OOS partners.

SECOORA researchers are developing a comprehensive data portal to provide discovery of, access to, and metadata about coastal ocean observations in the southeast US. Below are various ways to access the currently available data.

One project recently funded by SECOORA initiated development of species specific habitat models that integrate remotely sensed and in situ data to enhance stock assessments for species managed by the Council. The project during 2013/2014 was initiated to address red porgy, gray triggerfish, black seabass, and vermilion snapper. Gray triggerfish and red porgy are slated for assessment through SEDAR in 2014/15 and 2015/16 respectively.

#### *National Fish Habitat Plan and Southeast Aquatic Resource Partnership (SARP)*

In addition, the Council serves on the National Habitat Board and, as a member of the Southeast Aquatic Resource Partnership (SARP), has highlighted this collaboration by including the Southeast Aquatic Habitat Plan (SAHP) and associated watershed conservation restoration targets into the FEP. Many of the habitat, water quality, and water quantity conservation needs identified in the threats and recommendations Volume of the FEP are directly addressed by on-

the-ground projects supported by SARP. This cooperation results in funding fish habitat restoration and conservation intended to increase the viability of fish populations and fishing opportunity, which also meets the needs to conserve and manage Essential Fish Habitat for Council managed species or habitat important to their prey. To date, SARP has funded 53 projects in the region through this program. This work supports conservation objectives identified in the SAHP to improve, establish, or maintain riparian zones, water quality, watershed connectivity, sediment flows, bottoms and shorelines, and fish passage, and addresses other key factors associated with the loss and degradation of fish habitats. SARP also developed the Southern Instream Flow Network (SIFN) to address the impacts of flow alterations in the Southeastern US aquatic ecosystems which leverages policy, technical experience, and scientific resources among partners based in 15 states. Maintaining appropriate flow into South Atlantic estuarine systems to support healthy inshore habitats essential to Council managed species is a major regional concern and efforts of SARP through SIFN are envisioned to enhance state and local partners ability to maintain appropriate flow rates.

#### *Governor's South Atlantic Alliance (GSAA)*

Initially discussed as a South Atlantic Eco-regional Compact, the Council has also cooperated with South Atlantic States in the formation of a Governor's South Atlantic Alliance (GSAA). This will also provide regional guidance and resources that will address State and Council broader habitat and ecosystem conservation goals. The GSAA was initiated in 2006. An Executive Planning Team (EPT), by the end of 2007, had created a framework for the Governors South Atlantic Alliance. The formal agreement between the four states (NC, SC, GA, and FL) was executed in May 2009. The Agreement specifies that the Alliance will prepare a "Governors South Atlantic Alliance Action Plan" which will be reviewed annually for progress and updated every five years for relevance of content. The Alliance's mission and purpose is to promote collaboration among the four states, and with the support and interaction of federal agencies, academe, regional organizations, non-governmental organizations, and the private sector, to sustain and enhance the region's coastal and marine resources. The Alliance proposes to regionally implement science-based actions and policies that balance coastal and marine ecosystems capacities to support both human and natural systems. The GSAA Action Plan was released in December 2010 and describes the four Priority Issue Areas that were identified by the Governors to be of mutual importance to the sustainability of the region's resources: Healthy Ecosystems; Working Waterfronts; Clean Coastal and Ocean Waters; and Disaster-Resilient Communities. The goals, objectives, actions, and implementation steps for each of these priorities were further described in the GSAA Implementation Plan released in July 2011. The final Action Plan was released on December 1, 2010 and marked the beginning of intensive work by the Alliance Issue Area Technical Teams (IATTs) to develop implementation steps for the actions and objectives. The GSAA Implementation Plan was published July 6, 2011, and the Alliance has been working to implement the Plan through the IATTs and two NOAA-funded Projects. The Alliance also partners with other federal agencies, academia, non-profits, private industry, regional organizations, and others. The Alliance supports both national and state-level ocean and coastal policy by coordinating federal, state, and local entities to ensure the sustainability of the region's economic, cultural, and natural resources. The Alliance has organized itself around the founding principles outlined in the

GSAA Terms of Reference and detailed in the GSAA Business Plan. A team of natural resource managers, scientists, and information management system experts have partnered to develop a Regional Information Management System (RIMS) and recommend decision support tools that will support regional collaboration and decision-making. In addition to regional-level stakeholders, state and local coastal managers and decision makers will also be served by this project, which will enable ready access to new and existing data and information. The collection and synthesis of spatial data into a suite of visualization tools is a critical step for long-term collaborative planning in the South Atlantic region for a wide range of coastal uses. The Council's Atlas presents the spatial representations of Essential Fish Habitat, managed areas, regional fish and fish habitat distribution, and fishery operation information and it can be linked to or drawn on as a critical part of the collaboration with the RIMS.

### *South Atlantic Landscape Conservation Cooperative*

One of the more recent collaborations is the Council's participation as Steering Committee member for the newly establish South Atlantic Landscape Conservation Cooperative (SALCC). Landscape Conservation Cooperatives (LCCs) are applied conservation science partnerships focused on a defined geographic area that informs on-the-ground strategic conservation efforts at landscape scales. LCC partners include DOI agencies, other federal agencies, states, tribes, non-governmental organizations, universities, and others. The newly formed Department of Interior Southeast Climate Services Center (CSC) has the LCCs in the region as their primary clients. One of the initial charges of the CSCs is to downscale climate models for use at finer scales.

The SALCC developed a Strategic Plan through an iterative process that began in December 2011. The plan provides a simple strategy for moving forward over the next few years. An operations plan was developed under direction from the SALCC Steering Committee to redouble efforts to develop version 1.0 of a shared conservation blueprint by spring-summer of 2014. The SALCC is developing the regional blueprint to address the rapid changes in the South Atlantic including but not limited to climate change, urban growth, and increasing human demands on resources which are reshaping the landscape. While these forces cut across political and jurisdictional boundaries, the conservation community does not have a consistent cross-boundary, cross-organization plan for how to respond. The South Atlantic Conservation Blueprint will be that plan. The blueprint is envisioned to be a spatially-explicit map depicting the places and actions need to sustain South Atlantic LCC objectives in the face of future change. The steps to creating the blueprint include development of: indicators and targets (shared metrics of success); the State of the South Atlantic (past, present, and future condition of indicators); and a Conservation Blueprint. Potential ways the blueprint could be used include: finding the best places for people and organizations to work together; raising new money to implement conservation actions; guiding infrastructure development (highways, wind, urban growth, etc.); creating incentives as an alternative to regulation; bringing a landscape perspective to local adaptation efforts; and locating places and actions to build resilience after major disasters (hurricanes, oil spills, etc.). Integration of connectivity, function, and threats to river, estuarine and marine systems supporting Council managed species is supported by the SALCC and enhanced by the Council being a voting member of

its Steering Committee. In addition, the Council's Regional Atlas presents spatial representations of Essential Fish Habitat, managed areas, regional fish and fish habitat distribution, and fishery operation information and it be linked to or drawn on as a critical part of the collaboration with the recently developed SALCC Conservation Planning Atlas.

### **Building Tools to support EBM in the South Atlantic Region**

The Council has developed a Habitat and Ecosystem Section of the website <http://www.safmc.net/ecosystem/Home/EcosystemHome/tabid/435/Default.aspx> and, in cooperation with the Florida Wildlife Research Institute (FWRI), developed a Habitat and Ecosystem Internet Map Server (IMS). The IMS was developed to support Council and regional partners' efforts in the transition to EBM. Other regional partners include NMFS Habitat Conservation, South Atlantic States, local management authorities, other Federal partners, universities, conservation organizations, and recreational and commercial fishermen. As technology and spatial information needs evolved, the distribution and use of GIS demands greater capabilities. The Council has continued its collaboration with FWRI in the now evolution to Web Services provided through the regional SAFMC Habitat and Ecosystem Atlas ([http://ocean.floridamarine.org/safmc\\_atlas/](http://ocean.floridamarine.org/safmc_atlas/)) and the SAFMC Digital Dashboard ([http://ocean.floridamarine.org/safmc\\_dashboard/](http://ocean.floridamarine.org/safmc_dashboard/)). The Atlas integrates services for the following:

Species distribution and spatial presentation of regional fishery independent data from the SEAMAP-SA, MARMAP, and NOAA SEFIS systems; SAFMC Fisheries: ([http://ocean.floridamarine.org/SA\\_Fisheries/](http://ocean.floridamarine.org/SA_Fisheries/))

Essential Fish Habitat and Essential Fish Habitat Areas of Particular Concern; SAFMC EFH: ([http://ocean.floridamarine.org/sa\\_efh/](http://ocean.floridamarine.org/sa_efh/))

Spatial presentation of managed areas in the region; SAFMC Managed Areas: ([http://ocean.floridamarine.org/safmc\\_managedareas/](http://ocean.floridamarine.org/safmc_managedareas/))

An online life history and habitat information system supporting Council managed, State managed, and other regional species was developed in cooperation with FWRI. The Ecospecies system is considered dynamic and presents, as developed, detailed individual species life history reports and provides an interactive online query capability for all species included in the system: <http://atoll.floridamarine.org/EcoSpecies>

#### Web Services System Updates:

- Essential Fish Habitat (EFH) – displays EFH and EFH-HAPCS for SAFMC managed species and NOAA Fisheries Highly Migratory Species.
- Fisheries - displays Marine Resources Monitoring, Assessment, and Prediction (MARMAP) and Southeast Area Monitoring and Assessment Program South Atlantic (SEAMAP-SA) data.
- Managed Areas - displays a variety of regulatory boundaries (SAFMC and Federal) or management boundaries within the SAFMC’s jurisdiction.
- Habitat – displays habitat data collected by SEADESC, Harbor Branch Oceanographic Institute (HBOI), and Ocean Exploration dives, as well as the SEAMAP shallow and ESDIM deepwater bottom mapping projects, multibeam imagery, and scientific cruise data.
- Multibeam Bathymetry - displays a variety of multibeam data sources and scanned bathymetry charts.
- Nautical Charts – displays coastal, general, and overview nautical charts for the SAFMC’s jurisdictional area.

#### **Ecosystem Based Action, Future Challenges and Needs**

The Council has implemented ecosystem-based principles through several existing fishery management actions including establishment of deepwater Marine Protected Areas for the Snapper Grouper fishery, proactive harvest control rules on species (e.g., dolphin and wahoo) which are not overfished, implementing extensive gear area closures which in most cases eliminate the impact of fishing gear on Essential Fish Habitat, and use of other spatial management tools including Special Management Zones. Pursuant to development of the Comprehensive Ecosystem-Based Amendment, the Council has taken an ecosystem approach to protect deepwater ecosystems while providing for traditional fisheries for the Golden Crab and Royal Red shrimp in areas where they do not impact deepwater coral habitat. The stakeholder based process taps in on an extensive regional Habitat and Ecosystem network. Support tools facilitate Council deliberations and with the help of regional partners, are being refined to address long-term ecosystem management needs.

One of the greatest challenges to the long-term move to EBM in the region is funding high priority research, including but not limited to, comprehensive benthic mapping and ecosystem model and management tool development. In addition, collecting detailed information on fishing fleet dynamics including defining fishing operation areas by species, species complex, and season, as well as catch relative to habitat is critical for assessment of fishery, community, and habitat impacts and for Council use in place based management measures. Additional resources need to be dedicated to expand regional coordination of modeling, mapping, characterization of species use of habitats, and full funding of regional fishery independent surveys (e.g., MARMAP, SEAMAP, and SEFIS) which are linking directly to addressing high priority management needs. Development of ecosystem information systems to support Council management should build on existing tools (e.g., Regional Habitat and Ecosystem GIS and Arc Services) and provide resources to regional cooperating partners for expansion to address long-term Council needs.



The FEP and CE-BA 1 complement, but do not replace, existing FMPs. In addition, the FEP serves as a source document to the CE-BAs. NOAA should support and build on the regional coordination efforts of the Council as it transitions to a broader management approach. Resources need to be provided to collect information necessary to update and refine our FEP and support future fishery actions including but not limited to completing one of the highest priority needs to support EBM, the completion of mapping of near-shore, mid-shelf, shelf edge, and deepwater habitats in the South Atlantic region. In developing future FEPs, the Council will draw on SAFEs (Stock Assessment and Fishery Evaluation reports) which NMFS is required to provide the Council for all FMPs implemented under the Magnuson-Stevens Act. The FEP, which has served as the source document for CE-BAs, could also meet some of the NMFS SAFE requirements if information is provided to the Council to update necessary sections.

### **EFH and EFH-HAPC Designations Translated to Cooperative Habitat Policy Development and Protection**

The Council actively comments on non-fishing projects or policies that may impact fish habitat. **Appendix A** of the Comprehensive Amendment Addressing Essential Fish Habitat in Fishery Management Plans of the South Atlantic Region (SAFMC 1998b) outlines the Council's comment and policy development process and the establishment of a four-state Habitat Advisory Panel. Members of the Habitat Advisory Panel serve as the Council's habitat contacts and professionals in the field. AP members bring projects to the Council's attention, draft comment letters, and attend public meetings. With guidance from the Advisory Panel, the Council has developed and approved policies on:

1. Energy exploration, development, transportation, and hydropower re-licensing;
2. Beach dredging and filling and large-scale coastal engineering;
3. Protection and enhancement of submerged aquatic vegetation;
4. Alterations to riverine, estuarine, and nearshore flows;
5. Marine aquaculture;
6. Marine Ecosystems and Non-Native and Invasive Species: and
7. Estuarine Ecosystems and Non-Native and Invasive Species.

NOAA Fisheries, State and other Federal agencies apply EFH and EFH-HAPC designations and protection policies in the day-to-day permit review process. The revision and updating of existing habitat policies and the development of new policies is being coordinated with core agency representatives on the Habitat and Coral Advisory Panels. Existing policies are included at the end of this Appendix.

The Habitat and Environmental Protection Advisory Panel, as part of their role in providing continued policy guidance to the Council, is during 2013/14, reviewing and proposing revisions and updates to the existing policy statements and developing new ones for Council consideration. The effort is intended to enhance the value of the statements and support cooperation and collaboration with NOAA Fisheries Habitat Conservation Division and State and Federal partners in better addressing the Congressional mandates to the Council associated with designation and conservation of EFH in the region.

### **South Atlantic Bight Ecopath Model**

The Council worked cooperatively with the University of British Columbia and the Sea Around Us project to develop a straw-man and preliminary food web models (Ecopath with Ecosim) to characterize the ecological relationships of South Atlantic species, including those managed by the Council. This effort was envisioned to help the Council and cooperators in identifying available information and data gaps while providing insight into ecosystem function. More importantly, the model development process provides a vehicle to identify research necessary to better define populations, fisheries, and their interrelationships. While individual efforts are still underway in the South Atlantic, only with significant investment of new resources through other programs will a comprehensive regional model be further developed.

The latest collaboration builds on the previous Ecopath model developed through the Sea Around Us project for the South Atlantic Bight with a focus on beginning a dialogue on the implications of potential changes in forage fish populations in the region that could be associated with environmental or climate change or changes in direct exploitation of those populations.

### **Essential Fish Habitat and Essential Fish Habitat Areas of Particular Concern**

Following is a summary of the current South Atlantic Council's EFH and EFH-HAPCs. Information supporting their designation was updated (pursuant to the EFH Final Rule) in the Council's Fishery Ecosystem Plan and Comprehensive Ecosystem Amendment:

#### **Snapper Grouper FMP**

Essential fish habitat for snapper grouper species includes coral reefs, live/hard bottom, submerged aquatic vegetation, artificial reefs, and medium to high profile outcroppings on and around the shelf break zone from shore to at least 600 feet (but to at least 2,000 feet for wreckfish) where the annual water temperature range is sufficiently warm to maintain adult populations of members of this largely tropical complex. EFH includes the spawning area in the water column above the adult habitat and the additional pelagic environment, including *Sargassum*, required for larval survival and growth up to and including settlement. In addition the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse snapper grouper larvae.

For specific life stages of estuarine dependent and nearshore snapper grouper species, essential fish habitat includes areas inshore of the 100-foot contour, such as attached macroalgae; submerged rooted vascular plants (seagrasses); estuarine emergent vegetated wetlands (saltmarshes, brackish marsh); tidal creeks; estuarine scrub/shrub (mangrove fringe); oyster reefs and shell banks; unconsolidated bottom (soft sediments); artificial reefs; and coral reefs and live/hard bottom.

Areas which meet the criteria for EFH-HAPCs for species in the snapper-grouper management unit include medium to high profile offshore hard bottoms where spawning normally occurs; localities of known or likely periodic spawning aggregations; nearshore hard bottom areas; The

Point, The Ten Fathom Ledge, and Big Rock (North Carolina); The Charleston Bump (South Carolina); mangrove habitat; seagrass habitat; oyster/shell habitat; all coastal inlets; all state-designated nursery habitats of particular importance to snapper grouper (e.g., Primary and Secondary Nursery Areas designated in North Carolina); pelagic and benthic *Sargassum*; Hoyt Hills for wreckfish; the *Oculina* Bank Habitat Area of Particular Concern; all hermatypic coral habitats and reefs; manganese outcroppings on the Blake Plateau; and Council-designated Artificial Reef Special Management Zones (SMZs). In addition, the Council through CEBA 2 (SAFMC 2011) designated the deepwater snapper grouper MPAs and golden tilefish and blueline tilefish habitat as EFH-HAPCs under the Snapper Grouper FMP as follows:

EFH-HAPCs for golden tilefish to include irregular bottom comprised of troughs and terraces inter-mingled with sand, mud, or shell hash bottom. Mud-clay bottoms in depths of 150-300 meters are HAPC. Golden tilefish are generally found in 80-540 meters, but most commonly found in 200-meter depths.

EFH-HAPC for blueline tilefish to include irregular bottom habitats along the shelf edge in 45-65 meters depth; shelf break or upper slope along the 100-fathom contour (150-225 meters); hardbottom habitats characterized as rock overhangs, rock outcrops, manganese-phosphorite rock slab formations, or rocky reefs in the South Atlantic Bight; and the Georgetown Hole (Charleston Lumps) off Georgetown, SC.

EFH-HAPCs for the snapper grouper complex to include the following deepwater Marine Protected Areas (MPAs) as designated in Snapper Grouper Amendment 14: Snowy Grouper Wreck MPA, Northern South Carolina MPA, Edisto MPA, Charleston Deep Artificial Reef MPA, Georgia MPA, North Florida MPA, St. Lucie Hump MPA, and East Hump MPA.

Deepwater Coral HAPCs designated in Comprehensive Ecosystem-Based Amendment 1 are designated as Snapper Grouper EFH-HAPCs: Cape Lookout Coral HAPC, Cape Fear Coral HAPC, Blake Ridge Diapir Coral HAPC, Stetson-Miami Terrace Coral HAPC, and Pourtalés Terrace Coral HAPC.

### **Shrimp FMP**

For penaeid shrimp, Essential Fish Habitat includes inshore estuarine nursery areas, offshore marine habitats used for spawning and growth to maturity, and all interconnecting water bodies as described in the Habitat Plan. Inshore nursery areas include tidal freshwater (palustrine), estuarine, and marine emergent wetlands (e.g., intertidal marshes); tidal palustrine forested areas; mangroves; tidal freshwater, estuarine, and marine submerged aquatic vegetation (e.g., seagrass); and subtidal and intertidal non-vegetated flats. This applies from North Carolina through the Florida Keys.

For rock shrimp, essential fish habitat consists of offshore terrigenous and biogenic sand bottom habitats from 18 to 182 meters in depth with highest concentrations occurring between 34 and 55 meters. This applies for all areas from North Carolina through the Florida Keys. Essential fish habitat includes the shelf current systems near Cape Canaveral, Florida, which provide

major transport mechanisms affecting planktonic larval rock shrimp. These currents keep larvae on the Florida Shelf and may transport them inshore in spring. In addition, the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse rock shrimp larvae.

Essential fish habitat for royal red shrimp include the upper regions of the continental slope from 180 meters (590 feet) to about 730 meters (2,395 feet), with concentrations found at depths of between 250 meters (820 feet) and 475 meters (1,558 feet) over blue/black mud, sand, muddy sand, or white calcareous mud. In addition, the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse royal red shrimp larvae.

Areas which meet the criteria for EFH-HAPCs for penaeid shrimp include all coastal inlets, all state-designated nursery habitats of particular importance to shrimp (for example, in North Carolina this would include all Primary Nursery Areas and all Secondary Nursery Areas), and state-identified overwintering areas.

### **Coastal Migratory Pelagics FMP**

Essential fish habitat for coastal migratory pelagic species includes sandy shoals of capes and offshore bars, high profile rocky bottom, and barrier island ocean-side waters, from the surf to the shelf break zone, but from the Gulf Stream shoreward, including *Sargassum*. In addition, all coastal inlets and all state-designated nursery habitats of particular importance to coastal migratory pelagics (for example, in North Carolina this would include all Primary Nursery Areas and all Secondary Nursery Areas).

For Cobia essential fish habitat also includes high salinity bays, estuaries, and seagrass habitat. In addition, the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse coastal migratory pelagic larvae.

For king and Spanish mackerel and cobia essential fish habitat occurs in the South Atlantic and Mid-Atlantic Bights.

Areas which meet the criteria for EFH-HAPCs include sandy shoals of Capes Lookout, Cape Fear, and Cape Hatteras from shore to the ends of the respective shoals, but shoreward of the Gulf stream; The Point, The Ten-Fathom Ledge, and Big Rock (North Carolina); The Charleston Bump and Hurl Rocks (South Carolina); The Point off Jupiter Inlet (Florida); *Phragmatopoma* (worm reefs) reefs off the central east coast of Florida; nearshore hard bottom south of Cape Canaveral; The Hump off Islamorada, Florida; The Marathon Hump off Marathon, Florida; The “Wall” off of the Florida Keys; Pelagic *Sargassum*; and Atlantic coast estuaries with high numbers of Spanish mackerel and cobia based on abundance data from the ELMR Program. Estuaries meeting this criteria for Spanish mackerel include Bogue Sound and New River, North Carolina; Bogue Sound, North Carolina (Adults May-September salinity >30 ppt); and New River, North Carolina (Adults May-October salinity >30 ppt). For Cobia they include Broad River, South Carolina; and Broad River, South Carolina (Adults & juveniles May-July salinity >25ppt).

### **Golden Crab FMP**

Essential fish habitat for golden crab includes the U.S. Continental Shelf from Chesapeake Bay south through the Florida Straits (and into the Gulf of Mexico). In addition, the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse golden crab larvae. The detailed description of seven essential fish habitat types (a flat foraminiferan ooze habitat; distinct mounds, primarily of dead coral; ripple habitat; dunes; black pebble habitat; low outcrop; and soft-bioturbated habitat) for golden crab is provided in Wenner et al. (1987). There is insufficient knowledge of the biology of golden crabs to identify spawning and nursery areas and to identify HAPCs at this time. As information becomes available, the Council will evaluate such data and identify HAPCs as appropriate through the framework.

### **Spiny Lobster FMP**

Essential fish habitat for spiny lobster includes nearshore shelf/oceanic waters; shallow subtidal bottom; seagrass habitat; unconsolidated bottom (soft sediments); coral and live/hard bottom habitat; sponges; algal communities (*Laurencia*); and mangrove habitat (prop roots). In addition the Gulf Stream is an essential fish habitat because it provides a mechanism to disperse spiny lobster larvae.

Areas which meet the criteria for EFH-HAPCs for spiny lobster include Florida Bay, Biscayne Bay, Card Sound, and coral/hard bottom habitat from Jupiter Inlet, Florida through the Dry Tortugas, Florida.

### **Coral, Coral Reefs, and Live/Hard Bottom Habitats FMP**

Essential fish habitat for corals (stony corals, octocorals, and black corals) incorporate habitat for over 200 species. EFH for corals include the following:

- A. Essential fish habitat for hermatypic stony corals includes rough, hard, exposed, stable substrate from Palm Beach County south through the Florida reef tract in subtidal waters to 30 m depth; subtropical (15°-35° C), oligotrophic waters with high (30-35‰) salinity and turbidity levels sufficiently low enough to provide algal symbionts adequate sunlight penetration for photosynthesis. Ahermatypic stony corals are not light restricted and their essential fish habitat includes defined hard substrate in subtidal to outer shelf depths throughout the management area.
- B. Essential fish habitat for *Antipatharia* (black corals) includes rough, hard, exposed, stable substrate, offshore in high (30-35‰) salinity waters in depths exceeding 18 meters (54 feet), not restricted by light penetration on the outer shelf throughout the management area.
- C. Essential fish habitat for octocorals excepting the order Pennatulacea (sea pens and sea pansies) includes rough, hard, exposed, stable substrate in subtidal to outer shelf depths within a wide range of salinity and light penetration throughout

the management area.

- D. Essential fish habitat for Pennatulacea (sea pens and sea pansies) includes muddy, silty bottoms in subtidal to outer shelf depths within a wide range of salinity and light penetration.

Areas which meet the criteria for EFH-HAPCs for coral, coral reefs, and live/hard bottom include: The 10-Fathom Ledge, Big Rock, and The Point (North Carolina); Hurl Rocks and The Charleston Bump (South Carolina); Gray's Reef National Marine Sanctuary (Georgia); The *Phragmatopoma* (worm reefs) reefs off the central east coast of Florida; Oculina Banks off the east coast of Florida from Ft. Pierce to Cape Canaveral; nearshore (0-4 meters; 0-12 feet) hard bottom off the east coast of Florida from Cape Canaveral to Broward County); offshore (5-30 meter; 15-90 feet) hard bottom off the east coast of Florida from Palm Beach County to Fowey Rocks; Biscayne Bay, Florida; Biscayne National Park, Florida; and the Florida Keys National Marine Sanctuary. In addition, the Council through CEBA 2 (SAFMC 2011) designated the Deepwater Coral HAPCs as EFH-HAPCs under the Coral FMP as follows:

Deepwater Coral HAPCs designated in Comprehensive Ecosystem-Based Amendment 1 as Snapper Grouper EFH-HAPCs: Cape Lookout Coral HAPC, Cape Fear Coral HAPC, Blake Ridge Diapir Coral HAPC, Stetson-Miami Terrace Coral HAPC, and Poutalés Terrace Coral HAPC.

### **Dolphin and Wahoo FMP**

EFH for dolphin and wahoo is the Gulf Stream, Charleston Gyre, Florida Current, and pelagic *Sargassum*. This EFH definition for dolphin was approved by the Secretary of Commerce on June 3, 1999 as a part of the South Atlantic Council's Comprehensive Habitat Amendment (SAFMC 1998b) (dolphin was included within the Coastal Migratory Pelagics FMP at that time).

Areas which meet the criteria for EFH-HAPCs for dolphin and wahoo in the Atlantic include The Point, The Ten-Fathom Ledge, and Big Rock (North Carolina); The Charleston Bump and The Georgetown Hole (South Carolina); The Point off Jupiter Inlet (Florida); The Hump off Islamorada, Florida; The Marathon Hump off Marathon, Florida; The "Wall" off of the Florida Keys; and Pelagic *Sargassum*. This EFH-HAPC definition for dolphin was approved by the Secretary of Commerce on June 3, 1999 as a part of the South Atlantic Council's Comprehensive Habitat Amendment (dolphin was included within the Coastal Migratory Pelagics FMP at that time).

### **Pelagic *Sargassum* Habitat FMP**

The Council through CEBA 2 (SAFMC 2011) designated the top 10 meters of the water column in the South Atlantic EEZ bounded by the Gulfstream, as EFH for pelagic *Sargassum*.

## **Actions Implemented That Protect EFH and EFH-HAPCs**

### **Snapper Grouper FMP**

- Prohibited the use of the following gears to protect habitat: bottom longlines in the EEZ inside of 50 fathoms or anywhere south of St. Lucie Inlet, Florida; bottom longlines in the wreckfish fishery; fish traps; bottom tending (roller- rig) trawls on live bottom habitat; and entanglement gear.
- Established the *Oculina* Experimental Closed Area where the harvest or possession of all species in the snapper grouper complex is prohibited.
- Established deepwater Marine Protected Areas (MPAs) as designated in Snapper Grouper Amendment 14: Snowy Grouper Wreck MPA, Northern South Carolina MPA, Edisto MPA, Charleston Deep Artificial Reef MPA, Georgia MPA, North Florida MPA, St. Lucie Hump MPA, and East Hump MPA.

### **Shrimp FMP**

- Prohibition of rock shrimp trawling in a designated area around the *Oculina* Bank,
- Mandatory use of bycatch reduction devices in the penaeid shrimp fishery,
- Mandatory Vessel Monitoring System (VMS) in the Rock Shrimp Fishery.
- A mechanism that provides for the concurrent closure of the EEZ to penaeid shrimping if environmental conditions in state waters are such that the overwintering spawning stock is severely depleted.

### **Pelagic Sargassum Habitat FMP**

- Prohibited all harvest and possession of *Sargassum* from the South Atlantic EEZ south of the latitude line representing the North Carolina/South Carolina border (34° North Latitude).
- Prohibited all harvest of *Sargassum* from the South Atlantic EEZ within 100 miles of shore between the 34° North Latitude line and the Latitude line representing the North Carolina/Virginia border.
- Harvest of *Sargassum* from the South Atlantic EEZ is limited to the months of November through June.
- Established an annual Total Allowable Catch (TAC) of 5,000 pounds landed wet weight.
- Required that an official observer be present on each *Sargassum* harvesting trip. Require that nets used to harvest *Sargassum* be constructed of four inch stretch mesh or larger fitted to a frame no larger than 4 feet by 6 feet.

### **Coastal Migratory Pelagics FMP**

- Prohibited of the use of drift gillnets in the coastal migratory pelagic fishery.

### **Golden Crab FMP**

- In the northern zone, golden crab traps can only be deployed in waters deeper than 900 feet; in the middle and southern zones traps can only be deployed in waters deeper than

700 feet.

Northern zone - north of the 28°N. latitude to the North Carolina/Virginia border;

Middle zone - 28°N. latitude to 25° N. latitude; and

Southern zone - south of 25°N. latitude to the border between the South Atlantic and Gulf of Mexico Fishery Management Councils.

### **Coral, Coral Reefs and Live/Hard Bottom FMP**

- Established an optimum yield of zero and prohibiting all harvest or possession of these resources which serve as essential fish habitat to many managed species.
- Designated the *Oculina* Bank Habitat Area of Particular Concern.
- Expanded the *Oculina* Bank Habitat Area of Particular Concern (HAPC) to an area bounded to the west by 80°W. longitude, to the north by 28°30' N. latitude, to the south by 27°30' N. latitude, and to the east by the 100 fathom (600 feet) depth contour.
- Established the following two Satellite *Oculina* HAPCs: (1) Satellite *Oculina* HAPC #1 is bounded on the north by 28°30'N. latitude, on the south by 28°29'N. latitude, on the east by 80°W. longitude, and on the west by 80°3'W. longitude; and (2) Satellite *Oculina* HAPC #2 is bounded on the north by 28°17'N. latitude, on the south by 28°16'N. latitude, on the east by 80°W. longitude, and on the west by 80°3'W. longitude.
- Prohibited the use of all bottom tending fishing gear and fishing vessels from anchoring or using grapples in the *Oculina* Bank HAPC.
- Established a framework procedure to modify or establish Coral HAPCs.
- Established the following five deepwater CHAPCs:
  - Cape Lookout Lophelia Banks CHAPC;
  - Cape Fear Lophelia Banks CHAPC;
  - Stetson Reefs, Savannah and East Florida Lithoherms, and Miami Terrace (Stetson- Miami Terrace) CHAPC;
  - Pourtales Terrace CHAPC; and
  - Blake Ridge Diapir Methane Seep CHAPC.
- Within the deepwater CHAPCs, the possession of coral species and the use of all bottom damaging gear are prohibited including bottom longline, trawl (bottom and mid-water), dredge, pot or trap, or the use of an anchor, anchor and chain, or grapple and chain by all fishing vessels.



## **South Atlantic Council Policies for Protection and Restoration of Essential Fish**

### **Habitat**

#### **SAFMC Habitat and Environmental Protection Policy**

In recognizing that species are dependent on the quantity and quality of their essential habitats, it is the policy of the SAFMC to protect, restore, and develop habitats upon which fisheries species depend; to increase the extent of their distribution and abundance; and to improve their productive capacity for the benefit of present and future generations. For purposes of this policy, “habitat” is defined as the physical, chemical, and biological parameters that are necessary for continued productivity of the species that is being managed. The objectives of the SAFMC policy will be accomplished through the recommendation of no net loss or significant environmental degradation of existing habitat. A long-term objective is to support and promote a net-gain of fisheries habitat through the restoration and rehabilitation of the productive capacity of habitats that have been degraded, and the creation and development of productive habitats where increased fishery production is probable. The SAFMC will pursue these goals at state, Federal, and local levels. The Council shall assume an aggressive role in the protection and enhancement of habitats important to fishery species, and shall actively enter Federal, decision making processes where proposed actions may otherwise compromise the productivity of fishery resources of concern to the Council.

#### **SAFMC EFH Policy Statements**

In addition to implementing regulations to protect habitat from fishing related degradation, the Council in cooperation with NOAA Fisheries, actively comments on non-fishing projects or policies that may impact fish habitat. The Council adopted a habitat policy and procedure document that established a four-state Habitat Advisory Panel and adopted a comment and policy development process. Members of the Habitat Advisory Panel serve as the Council’s habitat contacts and professionals in the field. With guidance from the Advisory Panel, the Council has developed and approved a number of habitat policy statements which are available on the Habitat and Ecosystem section of the Council website (<http://www.safmc.net/ecosystem/Home/EcosystemHome/tabid/435/Default.aspx> ).

#### **References:**

SAFMC (South Atlantic Fishery Management Council). 1998a. Habitat Plan for the South Atlantic Region. South Atlantic Fishery Management Council, 1 Southpark Cir., Ste 306, Charleston, S.C. 29407-4699.

SAFMC (South Atlantic Fishery Management Council). 1998b. Comprehensive Amendment Addressing Essential Fish Habitat in Fishery Management Plans of the South Atlantic Region. South Atlantic Fishery Management Council, 1 Southpark Cir., Suite 306, Charleston, S.C. 29407-4699.

SAFMC (South Atlantic Fishery Management Council). 2009a. Fishery Ecosystem Plan for the South Atlantic Region. South Atlantic Fishery Management Council, 4055 Faber Place, Ste 201, North Charleston, S.C. 29405.

SAFMC (South Atlantic Fishery Management Council). 2009b. Comprehensive Ecosystem-Based Amendment 1 for the South Atlantic Region. South Atlantic Fishery Management Council, 4055 Faber Place Drive, Suite 201; North Charleston, SC 29405.

SAFMC (South Atlantic Fishery Management Council). 2011. Comprehensive Ecosystem-Based Amendment 2 for the South Atlantic Region. South Atlantic Fishery Management Council, 4055 Faber Place Drive, Suite 201; North Charleston, SC 29405.

Wenner, E. L., G. F. Ulrich, and J. B. Wise. 1987. Exploration for golden crab, *Geryon fenneri*, in the south Atlantic Bight: distribution, population structure, and gear assessment. *Fishery Bulletin* 85:547-560.